

Poster Presentation

MS83.P13


Synthesis, spectroscopic and structural analysis of a melamine copper complex

A. DIREM¹, N. DADDA², W. FALEK¹, Z. BOUTOBBA¹, N. BENALI-CHERIF¹

¹Université "Abbes Laghrour", Laboratoire des Structures, Propriétés et Interactions Inter-Atomiques LASPI2A, Khencela, Algérie, ²Université d'Annaba, Faculté des Sciences, Département de Chimie, Algérie.

Owing to its multiple sites acting as H-bonds' donors, melamine and its derivatives are considered as excellent building blocks for the construction of various crystalline supramolecular architectures based on phosphate, sulfate, chloride and carboxylic anions [1-10] on one hand and chlorometallate anions on the other hand [11-13]. In our contribution, we will discuss the synthesis, FTIR spectra and the structural investigation using the single crystal X-ray diffraction of (H₂Melamine) [CuCl₅]Cl, (I). The structural analysis of (I) revealed that it consists of alternating negative and positive layers running through the a-axis direction. The negative layers are built up of alternatively CuCl₅³⁻ and discrete non-coordination Cl⁻ anions, whereas the positive sheets are formed of [H₂melamine²⁺]₂ Hydrogen-bonded dimers. The N-H...N and N-H...Cl interactions are assembled to build three-dimensional H-bond patterns which insure the cohesion within the lattice (Fig. 01). A previous work showed that (I) exhibits antiferromagnetic properties [14]

[1] (a) D. J. A. De Ridder, K. Goubitz, V. Brodski, R. Peschar & H. Schenk. (2004). *Helv. Chim. Acta*, **87**. 1894. (b) V. Brodski, R. Peschar, H. Schenk, A. Brinkmann, E. R. H. van Eck, A. P. M. Kentgens, B. Coussens & A. Braam. (2004). *J. Phys. Chem. B*, **108**. 150, [2] (f) X. L. Zhang & X. M. Chen. (2005). *Cryst. Growth Des.*, **5**. 617. (g) X. L. Zhang, B. H. Ye & X. M. Chen. (2005). *Cryst. Growth Des.*, **5**. 1609. (h) J. P. Mathias, E. E. Simanek, J. A. Zerkowski, C. T. Seto & G. M. Whitesides. (1994). *J. Am. Chem. Soc.*, **116**, [3] (l) B. Dolling, A. G. Orpen, J. Starbuck & X. M. Wang. (2001). *Chem. Commun.*, 567. (m) C. J. Adams, H. M. Colquhoun, P. C. Crawford, M. Lusi & A. G. Orpen. (2007). *Angew. Chem. Int. Ed.*, **46**. 1124. (n) D. F. Weng, B. W. Wang, Z. M. Wang & S. Gao. (2011). Cr



Keywords: copper complex, single crystal X-ray diffraction, H-bonds